



- If the American public embraced a healthier lifestyle:
 - no smoking, no or moderate alcohol consumption
 - limited or no exposure to toxic chemicals
 - healthy nutrition
 - balance of exercise and rest
 - stress management
 - social integration


93% of diabetes, 81% of heart attacks, 50% of strokes, and 36% of all cancers could be prevented.

Ford, et al Arch Intem Med. 2009; 169(15):1355-62.









- CAD is leading cause of death in United States.
- CAD was considered primarily a lipid accumulation mediated disease; now shown to involve an ongoing inflammatory response. Christodoulidis G, et al. Cardiol Rev 2014 Jan 15

Copyright © 2009-2015 Integrative Medicine Concepts, LLC. All Rights Reserved

Periodontitis and Heart Disease



- Periodontitis recognized as an inflammatory disease of bacterial origin by American Academy of Periodontology.
- Plausibility of periodontal bacteria influencing vascular inflammation strengthened when *Porphyromonas gingivalis* isolated from human atheromatous plaques.
 Friedewald VE, et al. J Periodontol 2009; 80:1021-1032

Oral Health & Systemic Disease

- In recent years, periodontal diseases have been associated with systemic diseases such as rheumatoid arthritis, cardiovascular-disease, diabetes mellitus, chronic respiratory diseases and adverse pregnancy outcomes including preterm low-birth weight (PLBW) and pre-eclampsia.
- Improving periodontal health before or during pregnancy may prevent or reduce the occurrences of these adverse pregnancy outcomes and, therefore, reduce the maternal and perinatal morbidity and mortality. Parihar AS, et al. J Int Oral Health 2015; Jul;7(7):137-42.

Copyright © 2009-2015 Integrative Medicine Concepts, LLC. All Rights Reserved

Periodontitis and Cancer?


- Almost 60,000 Americans develop head and neck cancer annually and 12,000 dye from the disease.
- Chronic periodontitis may be independently associated with HNSCC through direct toxic effects of bacteria and their products, and/or through indirect effects of inflammation.
- It may also facilitate the acquisition and persistence of oral HPV infection, a recently recognized risk factor for HNSCC. Han YW, et al. Adv Dent Res 2014; 26(1):47-55

Copyright © 2009-2015 Integrative Medicine Concepts, LLC. All Rights Reserved.

Inflammation and Cancer

- Activation of NF-κβ is vital for our body's ability to defend itself against infection and is a central mediator of inflammation.
- Many things activate NF-κβ and when it is persistently "turned on" it induces genes that are responsible for increasing cell survival, proliferation, migration and invasion. It also turns down growth regulators (e.g., p53) that suppress tumor growth.

Pal S, et al. J Inflamm 2014; Aug 9; 11:23



Osteoporosis




- Growing evidence osteoporosis may be driven, in part, by chronic low grade inflammation.
- Pro-inflammatory cytokines (IL-1, IL-6, TNF-alpha), accelerate bone loss via activation of osteoclasts, inhibit collagen production in osteoblasts and enhance breakdown of the extracellular matrix.
- Large study of older women found almost a 50% increased risk of hip fracture in those with highest levels of inflammatory markers.

Barbour KE, et al. J Bone Miner Res 2014; 29(9):2057-64 Copyright © 2009-2015 Integrative Medicine Concepts, LLC. All Rights Reserved.

Factors That Drive Inflammation and Insulin Resistance

- Pattern of central obesity
- Western-dietary pattern, high fructose
- Sedentary lifestyle, lack of exercise
- Prolonged psychosocial stress
- Alterations in gut flora and intestinal permeability
- Environmental exposures (smoke, toxins)


Copyright © 2009-2015 Integrative Medicine Concepts, LLC. All Rights Reserved.




Obesity and Insulin Growth Factors

- Obesity associated with multiple factors that increase risk for heart disease, diabetes and certain cancers such as inflammation, insulin resistance, high blood sugar and insulin-growth factors (IGF).
- Elevated insulin increases tumor growth and aggressiveness.
- IGF-1 and IGF-2 have been identified as tumor promoters in multiple studies.

Belardi V, et al. J Mammary Gland Biol Neoplasia 2013; 18(3-4):277-89





Mediterranean Diet



• Traditional Mediterranean dietary pattern has shown superior anti-inflammatory effects when compared with typical North American and Northern European dietary patterns in most observational and interventional studies.

Galland L. Nutr Clin Pract 2010; Dec;25(6):634-40.

Copyright © 2009-2015 Integrative Medicine Concepts, LLC. All Rights Reserved

 Study > 2000 women confirmed "Western" diet increased breast cancer risk, while Med Diet lowered risk of all *breast cancer*




subtypes, particularly triple-negative tumors.

- Review found strong evidence of a beneficial role of the Mediterranean diet on *oral and pharyngeal cancer*.
- High adherence associated with significant reduction in overall cancer mortality (10%), colorectal cancer (14%), prostate cancer (4%) and aerodigestive cancer (56%).

Castello A, et al. Br J Cancer 2014; 111(7):1454-62 Filomeno M, et al. Br J Cancer 2014; 111(5):981-6. Schwingshacki L, et al. Int J Cancer 2014; 135(8):1884-97.








Non-alcoholic Fatty Liver Disease

- Small study found significant reductions in systolic blood pressure, body fat, liver enzymes (P = .04) and insulin resistance in children with NAFLD at 3 and 6 months (P < .05) on a *low fructose, low GL diet.*
- Four week study found low fructose diet in Hispanic adolescents increased insulin sensitivity and lowered hs-CRP.

Mager DR, et al. JPEN J Parenter Enteral Nutr 2013; Aug 23. Jin R, et al. Nutrients 2014; 6:3187–3201








Resolution Biology

- Inflammation always considered a passive process.
- Newer research shows that resolution of selflimited acute inflammation is an active, programmed response.
- Omega 3 fatty acids produce specialized proresolving mediators (SPM) – resolvins, protectins and maresins.



DHA + Aspirin and Periodontitis

- Harvard/Beth Israel study: 55 adults with moderate periodontitis randomized to 2,000 mg of DHA or identical placebo plus 81 mg/d of aspirin.
- DHA supplementation decreased mean pocket depth and gingival index. Significant adjusted differences were found between DHA and control for both gingival crevicular fluid hs-CRP (-5.3 ng/mL) and IL- 1β (p=.02) but systemic hs-CRP, plaque index or bleeding on probing.

Naqvi AZ, et al. J Dent Res 2014; 93 (8):767-773. Copyright © 2009-20151nteg nuive Medicine Concepts, LLC. All Rights Reserved.

Cardiovascular Health



Population studies and intervention trials show omega-3 fatty acids protect against cardiovascular morbidity and mortality.

Systematic reviews of clinical trials provide contradictory results for *secondary* prevention of cardiovascular disease.

Smith DA. Ann Intern Med 2012 Dec 18;157(12):JC6-5

Copyright \otimes 2009-2015 Integrative Medicine Concepts, LLC. All Rights Reserved

Canadians and Omega 3

- The Omega-3 Index indicates the percentage of EPA+DHA in red blood cell fatty acids.
- Canadian government found that the mean Omega-3 Index level of Canadians aged 20-79 was 4.5%.
 - Levels higher for women, older adults, Asians and other non-white Canadians, omega-3 supplement users, and fish consumers; levels lower for smokers and people who were obese.
- Fewer than 3% of adults had levels associated with low CHD risk; 43% had levels associated with high risk.

Langlois K, et al. Health Rep 2015; Nov 18;26(11):3-11


Copyright © 2009-2015 Integrative Medicine Concepts, LLC. All Rights Reserved

Pregnancy



- Maternal DHA intake during pregnancy and lactation can prolong high risk pregnancies, increase birth weight, head circumference; enhance visual acuity, hand and eye co-ordination, attention, and information processing.
- Maternal supplementation of omega 3 during pregnancy significantly reduced risk of childhood asthma and food allergies.


Morse NL. Nutrients 2012; 4(7): 799-840 Kiemens CM, et al. BJOG 2011; 118(8):916-25



Calculations are bas	ed on 4 oz. serving s	ize		
CATEGORY	SPECIES	WEEKLY MERCURY	SUSTAINABILITY	CONCLUSION
CC EWG'S	SALMON	NaN%	<u>Best Choice:</u> Wild Alaska	ï⊗î
BEST BETS! Very High Omega-3s, Low Mercury, Sustainable	SARDINES	NaN%	<u>Best Choice:</u> Pacific	789 1
	MUSSELS	NaN%	<u>Best Choice:</u> Farmed	۳×۹
	RAINBOW TROUT	NaN%	<u>Best Choice:</u> Farmed	۳×۹
	ATLANTIC MACKEREL	NaN%	Best Choice: Not Trawled	۳×۲

Different Types of Fish Oil Supplementation is an alternative to eating fish; however, all supplements are not equal. Randomized, crossover study of 35 healthy individuals compared four popular brands/types of omega 3 fatty acids: Concentrated triglyceride (rTG) Hyl ester (EE) Phospholipid krill oil (PL). Latlav M, et al Lipids Health Ds 2014; 13:92

ГRT	Product	EPA & DHA per capsule*	Tested values	Label use: caps/day	Daily dosage of EPA + DHA
rTG	Nordic Naturals ProOmega®	325 mg EPA	329.6 mg EPA	2	EPA: 650 mg
	Triglyceride	225 mg DHA	226.0 mg DHA		DHA: 450 mg
EE	Minami MorEPA®	756 mg EPA	774.2 mg EPA	1	EPA: 756 mg
	Platinum Ethyl Ester	228 mg DHA	233.7 mg DHA		DHA: 228 mg
PL	Source Naturals ArcticPure®	75 mg EPA	78.0 mg EPA	2	EPA: 150 mg
	Krill Oil Phospholipid	45 mg DHA.	46.7 mg DHA.		DHA: 90 mg
TG	New Chapter				
	Wholemega [®] Salmon	90 mg EPA	96.4 mg EPA	2	EPA: 180 mg
	Oil Triglyceride	110 mg DHA	109.5 mg DHA		DHA: 220 mg




Intestinal Permeability and Systemic Inflammation

- When lipopolysaccharide (LPS), endotoxin from membrane of gram negative bacteria in intestinal lumen, enters bloodstream it binds Toll-like receptor 4, inflammation is activated, changing insulin signaling and triggering inflammatory mediators
- Chronic exposure to LPS may contribute to weight gain and type 2 DM. Obese people and diabetics have increased plasma LPS.


Boroni Moreira AP, et al. Nutr Hosp 2012; 27(2):382-90

Copyright © 2009-2015 Integrative Medicine Concepts, LLC. All Rights Reserved.

Stress and The Gut



- Noradrenaline and adenosine released directly into gut mucosa from sympathetic nerve endings of ENS, causing mast cells to degranulate, releasing tryptase, histamine and other mediators.
- These neuromediators activate lymphocytes, stimulate epithelial chloride secretion and increase epithelial permeability.
- Subsequent movement of luminal bacteria into mucosa triggers further immune and inflammatory events.




Leaky Gut and Depression?


- LPS found to be significantly higher in patients with chronic depression than in those without.
- Multiple linear regression analysis revealed that TLR4 is an independent risk factor relating to severity of major depression.


Maes M, et al. J Affect Disord. 2011;11:414–418. Hung YY, et al. Psychiatry Res 2014; pii: S0165-1781(14)00653-2.


Copyright \oplus 2009-2015 Integrative Medicine Concepts, LLC. All Rights Reserved




Pre and Probiotics Regulate/modulate immune functions, reduce risk intestinal infection. Improve intestinal barrier functions, reduce metabolic endotoxemia Induce hypo-responsiveness to food antigens Improve glucose control and reduce inflammatory cytokines.











Absorption



- Most research uses 1200-2000 mg of *curcumin*
- Curcumin is poorly absorbed from the GI tract.
- Preparations that increase absorption include: nanoformulations, phospholipid, and those with piperine.
- Doses up to 12 grams per day for three months not associated with toxicity HOWEVER, GI upset is common complaint.


Kanai M. World J Gastroenterol 2014; 20(28):9384-91. Kakarala M, et al. Breast Cancer Res Treat 2009; 122:777-785

Copyright © 2009-201 5 Integrative Medicine Concepts, LLC. All Rights Reserved

	Amount Per Serving	% Daily Value*
Turmeric root extract 4:1	500 mg	•
Serving Size: 1 Tablet		
Serving Per Container: 60		
	Amount Per Serving	% Daily Value
Total Carbohydrate	<1 g	<1%†
Turmeric extract (root), 95% curcuminoids	450 mg	•
Serving Size:1 capeule	50 mg	*
Serving Size:1 capeule		
Serving Size:1 cappule Serving Per Container: 120	50 mg Amount Per Serving 500 mg	
Turmeric (root) Serving Size:1 capsule Serving Per Container: 120 Turmeric (curcuma longa) extract (root) (standardized to contain 95% curcuminoids) Bioperine black pepper (piper nigrum) extract (fluit) (standardized to contain 95% piperin	Amount Per Serving 500 mg	%Daily Value
Serving Size:1 capsule Serving Per Container: 120 Turmeric (curcuma longa) extract (root) (standardized to contain 95% curcuminoids)	Amount Per Serving 500 mg	%Daily Value •
Serving Size:1 capsule Serving Per Container: 120 Turmeric (curcuma longa) extract (root) (standardized to contain 95% curcuminoids) Bioperine black pepper (piper nigrum) extract (fruit) (standardized to contains 95% piperin	Amount Per Serving 500 mg	%Daily Value •
Serving Size:1 capsule Serving Per Container: 120 Turmeric (curcuma longa) extract (root) (standardized to contain 95% curcuminoids) Bioperine black pepper (piper nigrum) extract (fruit) (standardized to contains 95% piperin Supplement Facts	Amount Per Serving 500 mg	%Daily Value •
Serving Size:1 capsule Serving Per Container: 120 Turmeric (curcuma longa) extract (root) (standardized to contain 95% curcuminoids) Bioperine black pepper (piper nigrum) extract (fruit) (standardized to contains 95% piperin Supplement Facts Serving Size: Two Capsules	Amount Per Serving 500 mg	%Daily Value •

Green Tea (Camellia sinensis) White, green, oolong and black teas contain polyphenols and flavonoids, particularly catechins. Dominant catechin in green tea is epigallocatechin gallate (EGCG), which exerts potent anti-inflammatory and antioxidant activity.

Copyright © 2009-2015 Integrative Medicine Concepts, ILC. All Rights Reserved



Green Tea and Prostate Cancer



- RDBPCT 60 men with high-grade prostate intraepithelial neoplasia given 600 mg of EGCG or placebo for 1 year.
 - 3% in EGCG group converted to prostate cancer versus 30% in placebo group.
- UK study:199 men (~74 years age) localized prostate cancer, randomized to oral capsule containing pomegranate, green tea, broccoli, turmeric; or identical placebo for 6 months.
 - Median rise in PSA in supplement group was 14.7% as opposed to 78.5% in the placebo group (P=0.0008).

Bettuzzi S, et al. Cancer Res 2006; 66(2):1234-40 Thomas R, et al. Prostate Cancer Prostatic Dis 2014; 17(2):180-6

Copyright \otimes 2009-2015 Integrative Medicine Concepts, LLC. All Rights Reserved.

BENEFITS OF PHYSICAL ACTIVITY






- Maintain body weight and muscle tone.
- Resets stress response
- American Cancer Society, exercise may reduce risk of cancer by reducing insulin and insulin-like growth factors levels, associated with increased cell/tumor growth.
- Review of 73 studies found 25% average risk reduction for breast cancer amongst physically active women.

Lynch, et al. Recent Results Cancer Res 2011; 186:13-42

Copyright © 2009-2015 Integrative Medicine Concepts, LLC. All Rights Reserved.



How Prolonged Stress is Harmful

- Consequences of sustained high levels of stress hormones, principally cortisol, increases our risk for:
 - Obesity, heart disease, diabetes, irritable bowel syndrome, heartburn, low libido, infertility, hypothyroidism, infection, delayed healing, and even certain cancers.
- Scientists believe that prolonged elevation of stress hormones *may shorten our lifespan by 15 years.*

A Nation on the Edge?



- According to CDC, 11% of Americans 12 and older take anti-depressant medications, a 400% increase from the 1980s, ~203 million prescriptions in 2011
- In 2011, ~63 million prescriptions for Xanax and Ativan (Xanax #1 psychiatric medication.
- Prevalence anxiety disorders ages 9 17 is 13%.
- 1 in 8 adolescents suffer from depression.

NCHS Data Brief Number 76, October 2011

Copyright © 2009-2015 Integrative Medicine Concepts, LLC. All Rights Reserved

Mind-Heart



• InterHeart Trial, an international study conducted in 52 countries that included more than 27,000 participants, revealed that depression, social isolation and a feeling of hopelessness increased a woman's risk of having a heart attack **more** than diabetes, high blood pressure, smoking, or obesity.

Rosengren A, et al. Lancet 2004; 364:953-962.


Copyright © 2009-2015 Integrative Medicine Concepts, LLC. All Rights Reserved.

Social Isolation and Health



- 148 studies found that social isolation is:
 - As bad as smoking 15 cigarettes a day.
 - As dangerous as being an alcoholic.
 - As harmful as never exercising.
 - Twice as dangerous as obesity.

Holt-Lunstad J, et al. PLoS Med 2010; 7:e1000316



<section-header><section-header><image><image>

Neuroplasticity

- Brain's ability to change its structure and function; regularly used neural circuits are strengthened and expanded, those rarely engaged are weakened and contract.
- Long-time meditators have greater activation of areas responsible for sustaining attention, processing empathy, integrating emotion and cognition, and perceiving the mental and emotional state of others.


Copyright © 2009-2015 Integrative Medicine Concepts, LLC. All Rights Reserved.

Mindfulness Meditation Systematic Review



- 47 trials (n=3515) found it improved:
 - Anxiety
 - Depression
 - Pain

Goyal M, et al. JAMA Intern Med 2014; 174(3):357-68



Concluding Thoughts

- Chronic inflammation and insulin resistance are believed to be the root cause of many chronic diseases. Obesity is a major driver.
- Nutrition and exercise are vitally important for reducing disease risk.
- Chronic, persistent stress is also a significant cause of chronic inflammation and insulin resistance. Learning how to enhance resiliency and